

Decline of the space density of quasars between $z = 2$ and $z = 4$

M. Vigotti

Istituto di Radioastronomia del CNR, Via P. Gobetti 101, I-40129 Bologna, Italy
 vigotti@ira.cnr.it

R. Carballo¹

*Dpto. de Matematica Aplicada y C. Computacion, Universidad de Cantabria, Avda. Los Castros s/n,
 E-39005 Santander, Spain*

C.R. Benn

Isaac Newton Group, Apartado 321, E-38700 Santa Cruz de La Palma, Spain

G. De Zotti

INAF-Oss. Astr. Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy

R. Fanti²

Dipartimento di Fisica, Università di Bologna, Via Irnerio 46, I-40126 Bologna, Italy

J.I. Gonzalez Serrano

Instituto de Fisica de Cantabria (CSIC-UC), Avda. Los Castros s/n, E-39005 Santander, Spain

K.-H. Mack^{3,4}

Istituto di Radioastronomia del CNR, Via P. Gobetti 101, I-40129 Bologna, Italy

J. Holt

Department of Physics and Astronomy, University of Sheffield, Hicks Building, Sheffield, S3 7RH, UK

ABSTRACT

We define a new complete sample of 13 optically-luminous radio quasars ($M_{AB}(1450\text{\AA}) < -26^m9$ and $\log P_{1.4\text{GHz}}(\text{WHz}^{-1}) > 25.7$) with redshift $3.8 < z < 4.5$, obtained by cross-correlating the FIRST radio survey and the APM catalogue of POSS-I. We measure the space density to be $(1.0 \pm 0.3) \text{ Gpc}^{-3}$, a factor 1.9 ± 0.7 smaller than the space density of similar quasars at $z \approx 2$ (FBQS). Using a new measurement of the radio-loud fraction of quasars we find that at $z = 4$ the total space density of quasars with $M_{AB}(1450\text{\AA}) < -26^m9$ is $(7.4 \pm 2.6) \text{ Gpc}^{-3}$. This is a factor 1.8 ± 0.8 less than the space density at $z \approx 2$, found by the 2dF quasar survey. This $z = 2 / z = 4$ ratio, consistent with that of the radio-loud quasars, is significantly different from the ratio ~ 10 found for samples including lower-luminosity quasars. This suggests that the decline of the space density beyond $z \approx 2$ is slower for optically-luminous quasars than for less-luminous ones.

Subject headings: cosmology: observations, early-universe, galaxies: evolution, galaxies: formation, quasars: general, surveys

1. Introduction

The space density of bright quasars is a factor ≈ 100 higher at $z = 2$ than at $z = 0$. Boyle et al. (2000; the 2dF survey) have shown that this increase can be described in terms of pure luminosity evolution, i.e. the shape of the luminosity function does not change with z , up to at least $z = 2.3$. At higher redshifts evidence for a decrease in space density, consistent with pure luminosity evolution, has been presented by several authors (e.g. Kenefick et al., 1995 and references therein, but see also Irwin et al. 1991). The Sloan Digital Sky Survey (SDSS, Fan et al. 2001a) has pushed exploration of the quasar phenomenon out to $z > 6$, the epoch of formation of the first massive black holes (and galaxies) (Becker et al. 2001b). The space density of quasars at $z \approx 4$ with $M_{AB}(1450 \text{ \AA}) < -25^m5$ ($\Omega = 1$ and $\Lambda = 0$, $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, as used throughout this paper) was found by Fan et al. (2001b) to be more than one order of magnitude lower than that found by Boyle et al. (2000) at $z \simeq 2.3$. Fan et al. (2001b) also found substantial deviations from pure luminosity evolution: the bright end of the quasar luminosity function at $z > 4$ appears to be flatter than found at lower redshifts, i.e. the decrease in space density with increasing redshift depends on luminosity.

Radio-loud quasars are a small subset of the quasar population, but as stressed by several authors (e.g. Peacock 1985; Dunlop & Peacock 1990; Shaver et al. 1996, 1999), selection in the radio drastically reduces contamination by foreground stars with optical colors similar to high-redshift quasars, and is unaffected by both intrinsic and extrinsic dust obscuration and reddening. Thus the high-redshift behaviour of the radio-loud quasar luminosity function is a robust probe of the abundance of massive objects at very early cosmological epochs. A second important motivation for radio searches of high- z quasars is understanding

the origin of the radio phenomenon. Differences, if any, in the early evolutionary behaviour of radio and optical activity would provide valuable clues.

Several searches for high-redshift radio-loud quasars have been carried out (e.g.: Shaver et al. 1996, 1999; Hook et al. 1998a; Jarvis & Rawlings 2000; Jarvis et al. 2001; Snellen et al. 2001; Hook et al., 2002). Shaver et al. (1996, 1999) obtained complete redshift information (i.e. no optical magnitude limit) for a sample of 442 radio-loud flat-spectrum quasars with flux densities $S_{2.7GHz} \geq 0.25 \text{ Jy}$. They argued for a drop by more than a factor of 10 in space density from $z \simeq 2.5$ to $z \simeq 6$, consistent with that found in the optical.

Hook et al. (1998b) reported indications that the peak in the quasar space density may have occurred at earlier cosmological epochs for more radio-luminous sources, the decline in space density with redshift being less pronounced for these radio sources. Jarvis & Rawlings (2000) analyzed the top decade in radio luminosity ($\log P_{2.7GHz}(\text{W Hz}^{-1}) > 28$, quasars and galaxies) of sources from the Parkes Half-Jansky Flat-Spectrum Sample. They deduce a decline by a factor ~ 4 between $z \simeq 2.5$ and $z \simeq 5$, less abrupt than envisaged by Shaver et al. (1996, 1999) and consistent with that measured by Dunlop & Peacock (1990) for radio sources in general. Jarvis et al. (2001) investigated the high-redshift evolution of the most-radio-luminous ($28.8 < \log P_{151MHz}(\text{W Hz}^{-1}) < 29.8$) steep-spectrum radio sources using three samples selected at low frequencies (3CRR, 6CE, and 6C*). They found the data consistent with a constant space density between $z \simeq 2.5$ and $z \simeq 4.5$ and exclude a decline as steep as suggested by Shaver et al. (1996, 1999). These results are not necessarily inconsistent, since different ranges of radio luminosity and different frequency selections were used.

No evidence for a rapid decline of the space density of soft-X-ray selected quasars at $z > 2.7$ was apparent in the ROSAT survey data analyzed by Miyaji et al. (2000).

In this paper we report the results of a search for $z \approx 4$ radio-loud quasars, down to radio flux densities thirty times fainter than in previous searches, and with no selection on radio spectral index. The paper is organized as follows:

¹Instituto de Física de Cantabria (CSIC-UC), Avda. Los Castros s/n, E-39005 Santander, Spain

²Istituto di Radioastronomia del CNR, Via Gobetti 101, I-40129 Bologna, Italy

³ASTRON, Postbus 2, NL-7990 AA Dwingeloo, The Netherlands

⁴Radioastronomisches Institut der Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

In Sect. 2 we describe the construction of our sample of $z = 4$ quasars.

In Sect. 3 we calculate the space density at $z \approx 4$, and compare it with that at $z \approx 2$ derived from a sample of quasars of similar optical and radio luminosity.

In Sect. 4 we calculate the $z \approx 4$ space density of all quasars optically brighter than $M_{AB}(1450\text{\AA}) < -26^m.9$, using a new determination of the fraction of radio loud quasars at that redshift, and compare it with that measured at $z = 2$.

Section 5 contains the discussion, and Sect. 6 our conclusions.

2. Sample of $z \approx 4$ radio quasars

2.1. Candidate selection

The broad-band optical color of quasars reddens dramatically with redshift, as the Lyman- α forest and the 912- $\text{\AA}$ Lyman-limit leave the blue band (Fig. 1). To obtain a sample of high-redshift quasars, we therefore sought optical identifications of FIRST radio sources (Becker et al. 1995, White et al. 1997) with objects catalogued by the Automated Plate Measuring facility (APM; (McMahon & Irwin 1992) as being star-like on the POSS-I E (red) plates, with red magnitude $E \leq 18^m.8$, and with measured color $O - E \geq 3$, or O fainter than the plate limit ($O \sim 21^m.7$).

The FIRST radio catalogue (version of July 2000) includes $\approx 660,000$ radio sources with peak flux density $S_{1.4\text{GHz}} > 1$ mJy in 2.24 sr and $7^h < RA < 17^h$, $-5^\circ < DEC < 57^\circ$, of which 2.138 sr are covered by the APM catalogue.

We sought optical identifications lying closer than $1''.5$ to each radio position, corresponding to ~ 3 standard deviations in the expected combined radio and optical positional errors (see Fig. 2).

Searching to a larger radius would have netted 1% more identifications, but the contamination from foreground stars rises rapidly with search radius (see again Fig. 2). A similar search radius ($1''.2$) was used by White et al (2000) in their search for First Bright Quasar Survey (FBQS) candidates. Our search yielded a starting list of 2323 candidates.

Trial spectroscopy showed that $\approx 85\%$ of these ‘stellar’ candidates are faint galaxies.

In order to reduce contamination by these




Fig. 1.— POSS-I $O - E$ color vs redshift for high-redshift quasars (a) with $3.6 \leq z \leq 4.5$, $E \leq 18^m.8$, from Djorgovski (2001, $z > 4$) and from NED (as of May 2002, $z < 4$); (b) the 18 high- z FIRST quasars identified in our search (Section 2.2). Filled symbols indicate measured $O - E$ colors, arrows are lower limits (e.g. > 3 for $E < 18^m.7$, O limit $21^m.7$). The quasars in (a) were selected in part on the basis of color, but nevertheless trace the typical reddening of a quasar with increasing z as the continuum redward of the $\text{Ly}\alpha$ line moves out of the observed O band (red limit ≈ 5000 $\text{\AA}$; prediction for typical quasar shown as solid curve, computed in a similar way as described in the Appendix). As redshift increases through $z \approx 4$, O -band shifts below the 912 $\text{\AA}$ Lyman-limit and the $O - E$ color rises rapidly. As redshift increases above 4.5, the predicted brightness of a quasar in E band drops sharply, so few are likely to be detected. The dashed line indicates $z = 3.8$, above which we believe selection by color to be nearly complete (Sect. 2.3).

galaxies, we ran the SExtractor program (Bertin & Arnouts 1996) on images of the 2323 candidates on the POSS-II red plates, which have better sensitivity and spatial resolution than POSS-I. Eighty-two percent of the candidates (1901 objects) were clearly extended (i.e. galaxies). Seventy-three candidates could not be found by SExtractor, either because they were defects on the POSS-I plate (62 objects) or due to confusion effects near a bright star (11 objects).

The remaining 349 candidates were sought in the Minnesota APS catalogue of POSS-I (Pennington et al. 1993). Fifty-five of the candidates, undetected by APM on the *O* plate (i.e. potentially very red), were detected by APS. These have $O - E < 3$, so were removed from the sample. This left 294 candidates, 194 of which we classified, on the basis of the SExtractor-generated parameters, as ‘star-like’, and 100 as ‘possibly star-like’. CCD images of a third half of these were obtained in $1''.3$ -seeing with the Loiano 1.5-m telescope of the Astronomical Observatory of Bologna. On these images, 25 of the 70 ‘star-like’, and 28 of the 33 ‘possibly star-like’ candidates were extended. We have not investigated further the 100 ‘possibly star-like’ objects. As a result we expect an incompleteness of $\approx 100 \times 5/33 = 15(\pm 4)$ stellar candidates.

2.2. Spectroscopy

Spectra were obtained of 121 of the 194 candidates, using the IDS spectrograph of the Isaac Newton Telescope (Benn et al. 2001, Holt et al., in prep.). Eighteen of the 121 (15%) are quasars with $3.6 < z < 4.5$. The fraction of high-redshift quasars varies little with the optical magnitude of the candidates, but depends strongly on radio flux density: $(7.7 \pm 2.9)\%$ for $S_{1.4GHz} < 10$ mJy and $(37 \pm 11)\%$ for $S_{1.4GHz} > 10$ mJy. Amongst the 73 candidates not observed spectroscopically, there are 6 with $S_{1.4GHz} > 10$ mJy and 67 with $S_{1.4GHz} < 10$ mJy. On the basis of the above high-redshift-quasar fractions, and accounting also for the ~ 15 missed stellar candidates (Sect. 2.1), we expect to have missed a total of 9 ± 2 high redshift quasars, i.e. the search is $\approx (67 \pm 5)\%$ complete or, equivalently, corresponds to an area on the sky of 1.432 sr.




Fig. 2.— The distributions of radio-optical separations r for the star-like candidates classified spectroscopically as quasars, galaxies and stars (total 194 objects). For quasars and galaxies, the dashed curves are of the form $r \cdot \exp(-r^2/2\sigma^2)$, with $\sigma = 0''.5$. For stars, the dashed straight line indicates the form expected for misidentifications with foreground stars.

2.3. Sample completeness

Our search for high-redshift quasars within this area will be incomplete for a number of other reasons, that we discuss below:

1) *Incompleteness of the APM catalogue.* According to McMahon et al. (2001), APM classifies correctly 92% of $E < 19^m$ objects on at least one of the O or E plates. Assuming that the APM E and O plate misclassifications are independent and occur at a similar rate, then a 92%-correct classification on at least one plate implies 72%-correct classification on individual plates. This last figure is a lower limit on the incompleteness of our E plate based star-like classification (the classification on the O and E plate cannot be completely independent).

For an independent estimate of our incompleteness, we searched for APM detections of 100 $z > 4$ quasars with $E \leq 19^m$ from the list of Djorgovski (2001), which were originally found either on the POSS-II plates or in SDSS (CCD based). All 100 were detected by APM. Eighteen of these quasars appeared extended in the APM E band, i.e. the search was $(82 \pm 4)\%$ complete.

As a further check we searched the APM catalogue for counterparts of 4949 objects with $16^m \leq E \leq 19^m$ catalogued as star-like in the APS. APM classifies as extended $(13 \pm 0.5)\%$ of these objects and misses a further $(3 \pm 0.3)\%$, i.e. is 84% complete. Because the APS classification algorithm (neural network) is believed to be more reliable than that of APM, we assume that disagreements about classification by APM and APS are due mainly to errors in the former.

To check the APS classification reliability, we constructed a sample of 190 quasars, 140 from the FBQS survey with $16^m < E < 18^m$ and 50 from the B3VLA QSS survey (Vigotti et al. 1997) with $18^m < E < 19^m$. Only 10 quasars were classified extended by APS, i.e. an incompleteness of 5%. This is three times smaller than the APM incompleteness, consistent with our assumption above.

The above two estimates of APM completeness are in good agreement and we assume a value of $(84 \pm 2)\%$. In addition, a few star-like objects might have been misclassified extended when we analysed the images with SExtractor, or at the stage of CCD imaging, but the number is likely to

be very small.

2) *Omission of quasars with extended radio emission.* Optical counterparts were sought at the positions of individual FIRST radio sources, but not at the mid-points of possible double sources. In the First Bright Quasar Survey (FBQS) in the South Galactic Cap, Becker et al. (2001a) also sought counterparts along the lines joining possible double sources of separation up to $30''$, and found that a search at the positions of individual radio components is 96% complete. At $z \sim 4$, the rest-frame frequency of observation is even higher than for FBQS, favouring more flat-spectrum compact sources. We therefore expect our sample to be more complete, and we assume a completeness of $(98 \pm 1)\%$.

3) *Incompleteness due to errors in the radio-optical positions.* The distribution of the radio - optical separations of the quasars (Fig. 2) is consistent with a rms error of $0''.5$, implying that our search out to radius $1''.5$ is $\approx 99\%$ complete.

4) *Incompleteness due to color selection.* The fraction of $z > 3.8$ quasars with $O - E > 3^m$ can be estimated approximately from the color distribution in Fig. 1a (taken from the literature). 81 of the 85 quasars with $z > 3.8$ have $O - E > 3$, i.e. this color criterion finds $(95 \pm 1.5)\%$ of quasars with $z > 3.8$, but the efficiency falls rapidly with decreasing redshift. We therefore restrict our sample to $z > 3.8$.

Combining the above, the completeness of our search for $z > 3.8$ quasars in FIRST is $\approx (77 \pm 6)\%$ over the effective area (Sect. 2.2) of 1.432 sr.

5) *Incompleteness of the FIRST radio survey.* The FIRST catalog is severely incomplete at faint flux densities. From Prandoni et al. (2001) we estimate a completeness of $\approx 50\%$ for $S \leq 1.25$ mJy and of $\approx 75\%$ for $1.25 \text{ mJy} \leq S \leq 1.5$ mJy. This flux-density-dependent incompleteness is in addition to that discussed above.

2.4. The $z \approx 4$ radio-loud sample

The 13 radio-loud quasars with $z \geq 3.8$, the Q_{z4RL} sample, are listed in Tab. 1.

The E magnitudes were recalibrated using APS catalogue (rms error $0^m.2$, better than APM, because the APS photometry was calibrated on a

plate-by-plate basis) and corrected for Galactic extinction. After re-calibration the limiting magnitude of the sample is $E = 19^m$.

The epoch-1950 absolute magnitudes in the rest-frame UV continuum, $M_{AB}(1450 \text{ \AA})$, were obtained from the E magnitudes, using a template quasar spectrum (the procedure is described in the Appendix). The 13 quasars are optically very luminous, $M_{AB}(1450 \text{ \AA}) \leq -26^m9$.

The radio luminosities were calculated using a median spectral index $\alpha_R = -0.3$, rms = 0.4 ($S_\nu \propto \nu^\alpha$), and range $\log P_{1.4\text{GHz}} (\text{W Hz}^{-1}) \geq 25.7$. The median spectral index was obtained via 5-GHz observations of some sources using the Effelsberg radio telescope (Holt et al. 2003, in prep.).

As a check of our method of deriving $M_{AB}(1450 \text{ \AA})$, we made, for 8 of the quasars, independent measurements based on recent I -band photometry obtained during service time at the 1.0-m Jacobus Kapteyn Telescope in La Palma (Holt et al., in prep.). At $z = 4.2$, I band corresponds to rest-frame wavelength $\sim 1540 \text{ \AA}$, close enough to the reference wavelength $1450 \text{ \AA}$ that we can K-correct using the median optical spectral index $\alpha_{OPT} = -0.7$ from Fan et al. (2001c). For two of the 8 quasars, 0831+52 and 1423+22, we found a large difference between the M_{AB} values derived using these two methods, the I -photometry giving M_{AB} respectively dimmer by 0^m4 and brighter by 1^m0 . These differences are probably due to intrinsic changes in luminosity between epochs 1950 and 2002. For the remaining 6 quasars the mean difference is 0.0 ± 0.1 , suggesting that any systematic error in our M_{AB} magnitude scale is $\lesssim 0^m1$.

The mean radio and optical luminosities of the sample are: $\langle \log P_{1.4\text{GHz}} \rangle = 26.6 \pm 0.2$ and $\langle M_{AB} \rangle = -27^m6 \pm 0^m2$, the median values being 26.8 and -27^m5 respectively.

3. Decline of the space density of radio loud quasars between $z = 2$ and $z = 4$

3.1. Comparison sample of $z \approx 2$ radio quasars

For a sample of radio quasars with comparable radio and optical luminosities, but at $z \approx 2$, we draw on the 0.817-sr First Bright Quasar Survey (FBQS, White et al. 2000). FBQS uses O , E magnitudes recalibrated using APS, with a limiting

magnitude $E = 17^m8$. Of the original 1238 quasar candidates, 636 were confirmed quasars. No spectrum is available for 108 of the 1238 candidates, i.e. the spectroscopy is 91% complete, corresponding to an effective area on the sky of 0.743 sr.

Of the 45 FBQS quasars with $1.8 \leq z \leq 2.4$, 23 have luminosities in the range of the Q_{z4RL} sample, i.e. $\log P_{1.4\text{GHz}} \geq 25.7$ and $M_{AB}(1450 \text{ \AA}) \leq -26^m9$.

The mean radio and optical luminosities values are $\langle \log P_{1.4\text{GHz}} \rangle = 26.6 \pm 0.2$ and $\langle M_{AB} \rangle = -27^m5 \pm 0.1$. The median values are 26.4 and -27^m4 respectively (consistent, within the statistical noise, with those of the $z = 4$ sample). We thus have a lower-redshift sample directly comparable with our high-redshift one. Fig. 3 shows the distributions of the sources in radio and optical luminosity and redshift for each sample, for the Einstein – de Sitter cosmology and for the *concordance model* (see Sect. 3.2)

This sample (Tab. 2) will be referred as the Q_{z2RL} sample. The radio luminosities $P_{1.4\text{GHz}}$ were calculated assuming a radio spectral index $\alpha_R = -0.3$. This is the median spectral index obtained for a subsample with GB6 4.8-GHz flux densities (Gregory et al. 1996), and is similar to that of the $z \approx 4$ sample. The absolute magnitudes $M_{AB}(1450 \text{ \AA})$ have been calculated from the E magnitudes using a K-correction = $2.5 \log(1+z)^{(1+\alpha_{OPT})}$ and $\alpha_{OPT} = -0.5$, that of the FBQS composite spectrum (Brotherton et al. 2001).

As an independent check, we also calculated the absolute magnitudes using the quasar template (following a procedure similar to that used at $z = 4$, see Appendix), obtaining similar results, within $\pm 0^m1$.

This sample will suffer from some of the same incompletenesses as the Q_{z4RL} sample (Section 2.3). The search for star-like objects $E \leq 17^m8$ in the APM will be $\approx (96 \pm 1)\%$ complete. An additional 4% of objects will be missed due to not searching at the mid-points of possible double radio sources (Becker et al. 2001a). No losses are expected due to color selection, nor for incompleteness of the FIRST at faint flux densities (these sources are brighter than 3 mJy). In total, the Q_{z2RL} sample of quasars is expected to be $(92 \pm 1.4)\%$ complete over 0.743 sr.

Table 1: Properties of the $z = 4$ radio-loud sample

RA	Dec	E	$O - E$	z	$S_{1.4GHz}$	$\log P_{1.4GHz}$	M_{AB}	z_{max}	ρ
J2000					mJy	WHz $^{-1}$	(1450Å)		Gpc $^{-3}$
07 25 18.29	+37 05 17.9	18.7	> 3.4	4.330	26.6	27.11	-26.9	4.379	0.0517
07 47 11.16	+27 39 03.6	17.8	> 4.5	4.170	1.1	25.70	-27.6	4.366	0.0530
08 31 41.68	+52 45 17.1	15.8	3.9	3.870	1.3	25.72	-29.8	4.500	0.0432
08 39 46.11	+51 12 03.0	18.9	> 3.2	4.420	41.6	27.32	-27.3	4.438	0.0472
09 18 24.38	+06 36 53.5	19.0	> 3.0	4.180	26.5	27.09	-26.9	4.206	0.0731
09 41 19.36	+51 19 32.3	18.0	> 4.1	3.850	2.5	26.00	-27.6	4.470	0.0450
10 53 20.42	-00 16 50.4	17.8	> 3.6	4.289	13.8	26.82	-27.7	4.483	0.0441
10 57 56.27	+45 55 53.0	16.9	3.9	4.110	1.1	25.70	-28.5	4.350	0.0544
12 11 34.41	+32 26 15.8	18.2	> 3.0	4.097	3.7	26.21	-27.2	4.421	0.0484
13 09 40.61	+57 33 09.1	18.3	> 3.6	4.274	11.3	26.73	-27.1	4.406	0.0495
13 25 12.52	+11 23 29.9	18.6	> 2.9	4.400	71.1	27.55	-27.5	4.455	0.0460
14 23 08.19	+22 41 57.5	18.7	> 3.2	4.316	35.4	27.23	-26.9	4.377	0.0519
16 39 50.51	+43 40 03.6	17.7	3.9	3.990	25.2	27.03	-27.8	4.500	0.0432

Table 1: The columns give: (1 - 2) J2000 optical coordinates, (3) E magnitudes recalibrated using APS; (4) APM $O - E$ color; (5) spectroscopic redshift; (6) FIRST radio flux density (peak or integrated, see next column), (7) radio luminosity, calculated using the peak flux density for $S_{1.4GHz} < 10$ mJy and the integrated flux density for $S_{1.4GHz} > 10$ mJy, as recommended by Ivezić et al. (2002, see their Fig. 4), and K-corrected using the median radio spectral index $\alpha_R = -0.3$ ($S_\nu \sim \nu^\alpha$, as used throughout this paper), (8) absolute magnitude M_{AB} at rest-frame 1450 Å (see Appendix), (9) maximum redshift out to which the quasar could have been detected (i.e. with $E < 19^m0$, $S_{1.4GHz} > 1$ mJy), (10) contribution of this source to space density, assuming effective area 1.432 sr, but without correction for the incompletenesses discussed in Sect. 2.3 (see text).

3.2. The space density of radio-loud quasars at $z \approx 4$ and at $z \approx 2$

We have two shells of volume: one defined by $1.8 \leq z \leq 2.4$ (which has been chosen to yield a good statistical sample around $z = 2$) and the other by $3.8 \leq z \leq 4.5$ (determined by the fact that at z larger than 4.5 the predicted brightness of a quasar in E band drops sharply, so few are likely to be detected). For each of the two samples of quasars, the space density was calculated as follows. For each source we computed $z_{max,r}$ and $z_{max,o}$ the maximum redshifts at which the source would reach the radio flux-density limit or the optical magnitude limit of the sample.

In addition, for the $z = 4$ sample of quasars, we calculated the *minimum* redshift at which the quasar could have been seen, since the K-correction could in principle cause the quasar to dim with decreasing redshift (Fig. 6). For all the sources we found this minimum redshift to be < 3.8 .

The maximum redshift z_{max} at which the quasar could have been included in the sample is the minimum of $z_{max,r}$ and $z_{max,o}$.

Just two of the quasars in the $z = 4$ sample are *radio limited* ($z_{max,r} \leq z_{max,o}$), while in the $z = 2$ sample there are none. The uncertainty in $z_{max,r}$ due to the assumption of an average radio spectral index (-0.3) is ~ 0.1 and implies a negligible effect on the calculated space density ($< 2\%$).

The lower limit on redshift is the sample limit z_0 (1.8 for the $z \approx 2$ sample, 3.8 for the $z \approx 4$ sample). The space density contributed by each quasar is the inverse of the volume of a shell bounded by z_0 and z_{max} , and with surface area equal to the effective area covered by the sample (0.743 and 1.432 sr for the Q_{z2RL} and the Q_{z4RL} samples respectively). This volume is:

$$V = k \left(\left(1 - \frac{1}{\sqrt{(1+z_{max})}} \right)^3 - \left(1 - \frac{1}{\sqrt{(1+z_0)}} \right)^3 \right)$$

where $k = 575$ Gpc 3 per sr.

Summing the contributions of individual sources, and normalising by area (Tabs. 1, 2), we ob-

Table 2: Properties of the $z = 2$ radio-loud sample

RA	Dec	E	$O - E$	z	$S_{1.4GHz}$	$\log P_{1.4GHz}$	M_{AB}	z_{max}	ρ
J2000					mJy	WHz $^{-1}$	(1450Å)		Gpc $^{-3}$
07 29 28.47	+25 24 51.99	17.2	0.6	2.303	6.8	26.05	-27.4	2.400	0.0761
07 58 13.94	+26 24 18.75	17.1	0.7	1.915	13.9	26.22	-27.1	2.400	0.0761
08 04 13.67	+25 16 34.07	17.4	0.7	2.302	3.5	25.76	-27.2	2.400	0.0761
08 57 26.95	+33 13 17.24	17.1	0.3	2.338	21.3	26.56	-27.6	2.400	0.0761
09 04 44.33	+23 33 54.05	16.9	0.3	2.244	107.3	27.23	-27.7	2.400	0.0761
09 10 54.18	+37 59 15.11	17.2	0.0	2.158	265.9	27.59	-27.3	2.400	0.0761
10 12 11.43	+33 09 25.96	17.3	0.0	2.260	196.3	27.50	-27.3	2.400	0.0761
10 35 10.98	+35 10 19.51	17.3	-0.2	1.955	22.9	26.45	-26.9	2.400	0.0761
10 36 41.94	+25 02 36.63	16.9	0.7	2.004	118.5	27.18	-27.5	2.400	0.0761
10 54 27.15	+25 36 00.33	16.9	1.4	2.400	3.0	25.72	-27.8	2.400	0.0761
11 22 41.46	+30 35 34.88	16.4	0.4	1.809	10.0	26.03	-27.8	2.400	0.0761
12 25 27.38	+22 35 12.72	16.2	0.2	2.043	3.9	25.71	-28.2	2.400	0.0761
12 28 24.98	+31 28 37.74	15.5	0.4	2.199	323.8	27.69	-29.0	2.400	0.0761
13 24 22.53	+24 52 22.25	17.4	1.5	2.357	4.9	25.92	-27.3	2.400	0.0761
13 45 20.42	+32 41 12.59	17.6	0.3	2.253	14.6	26.36	-27.0	2.400	0.0761
14 15 28.47	+37 06 21.49	17.4	0.9	2.360	409.7	27.85	-27.3	2.400	0.0761
14 16 17.38	+26 49 06.20	17.7	0.8	2.297	2.9	25.68	-26.9	2.400	0.0761
14 21 08.72	+22 41 17.42	16.8	0.1	2.181	4.7	25.85	-27.8	2.400	0.0761
14 23 26.07	+32 52 20.34	16.3	0.2	1.903	8.7	26.01	-27.9	2.400	0.0761
14 32 43.32	+41 03 28.04	17.4	0.7	1.963	261.7	27.51	-26.9	2.400	0.0761
16 03 54.15	+30 02 08.88	17.4	0.7	2.026	54.2	26.85	-27.0	2.400	0.0761
16 34 12.77	+32 03 35.45	16.6	0.3	2.341	176.5	27.48	-28.1	2.400	0.0761
16 51 37.56	+40 02 18.71	16.9	0.3	2.316	43.9	26.86	-27.7	2.400	0.0761

Table 2: The columns are as for Tab. 1, except that in column: (4) $O - E$ is the APM color using APS recalibration; (8) $M_{AB}(1450\text{\AA})$ is calculated from the E magnitude (the O magnitude is closer in rest-frame wavelength, but may include Ly α in the wing of the filter response); (9) the maximum redshift out to which the quasar could have been detected is calculated assuming $E < 17^m8$, $S_{1.4GHz} > 1$ mJy; (10) effective area 0.743 sr is assumed.

tain space densities of 0.65 ± 0.18 , and 1.75 ± 0.36 , quasars per Gpc 3 at $z \approx 4.2$ and $z \approx 2$ respectively.

Correcting for completeness ($77 \pm 5\%$ for the Q_{z4RL} sample, multiplied by 60% for $S \leq 1.5$ mJy), ($92 \pm 1.4\%$ for the Q_{z2RL} sample) the space densities become:

$$\rho_{z4RL} = 0.99 \pm 0.28 \text{ Gpc}^{-3}$$

$$\rho_{z2RL} = 1.90 \pm 0.47 \text{ Gpc}^{-3}$$

with ratio:

$$\rho_{z2RL}/\rho_{z4RL} = 1.9 \pm 0.7.$$

In calculating the errors, we also took into account possible systematic shifts of 0^m1 in the magnitude scale.

Had we adopted the currently more fashionable

cosmological *concordance model* with $\Omega_\Lambda = 0.8$, $\Omega_{\text{matter}} = 0.2$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the ratio of the comoving volumes $V(z=4)/V(z=2)$ would have been higher by a factor ≈ 1.25 . The derived radio and optical luminosities change by different amounts in the two redshift bins. E.g. $M_{AB} = -26^m9$ in the Einstein – de Sitter cosmology becomes in the *concordance model* $M_{AB} = -27^m16$ at $z = 2$, and $M_{AB} = -27^m34$ at $z = 4$. The same applies to the radio luminosities. To maintain comparable luminosities in the two samples, eight of the $z = 2$ quasars have to be omitted. The density ratio ρ_{z2RL}/ρ_{z4RL} then decreases by a factor $\simeq 0.81$.

4. The space density of optically-bright quasars at $z \approx 4$

In the previous section we measured the space density of optically bright ($M_{AB}(1450\text{\AA}) < -26^m9$) radio-loud ($\log P_{1.4} > 25.7$) quasars at $z \approx 4$. At this redshift the total space density of all optically-bright quasars (i.e. radio-loud + radio-quiet) is still poorly known, although SDSS should soon yield an accurate value. Here we obtain it from the space density of the Q_{z4RL} sample and from an accurate new measurement of the fraction of optically-bright quasars with $\log P_{1.4} > 25.7$ (radio loud).⁵ We indicate this fraction as G_{RL} .

4.1. The radio/optical fraction of quasars at $z \approx 4$

From Djorgovski's (2001) list of $z > 4$ quasars we extracted a sample of 191 optically-selected quasars lying in the area of sky covered by the FIRST survey. For 22 of these we found a FIRST radio source within $3''$. The number of chance coincidences will be negligible (the number of candidates does not change if the search radius is increased to $10''$), i.e. all of these are radio-loud quasars.

We are interested in the luminosity range covered by the Q_{z4RL} sample, $M_{AB}(1450\text{\AA}) \leq -26^m9$. From the reported magnitudes, which are APS or APM POSS-I or POSS-II R, we computed $M_{AB}(1450\text{\AA})$ for all 191 objects, with estimated rms and systematic errors both $\leq 0^m3$. The magnitude uncertainties are unimportant for our purposes, because we will see that the fraction of radio loud quasars, G_{RL} , changes little over a broad range of optical luminosity. Applying the correction factor for the incompleteness of the FIRST (see Sect. 2.3) we obtain:

$$G_{RL}(M_{AB}(1450\text{\AA}) < -26^m9) = (13.3 \pm 4.8)\%$$

$$G_{RL}(M_{AB}(1450\text{\AA}) > -26^m9) = (13.4 \pm 3.7)\%$$

i.e. there is no evidence that G_{RL} depends on optical luminosity. The mean value is:

$$G_{RL}(-25^m \gtrsim M_{AB}(1450\text{\AA}) \gtrsim -29^m) =$$

$$(13.4 \pm 3.0)\%.$$

This is consistent with the less accurate estimate $(12 \pm 6)\%$ by Stern et al. (2000).

In the next paragraph we will use this ratio and the space density of the Q_{z4RL} loud sample to find the space density of *all quasars* at $z \approx 4$ with $M_{AB} < -26^m9$.

4.2. The luminosity function of optically bright quasars

Above we measured a radio-loud fraction $G_{RL} = (13.4 \pm 3.0)\%$ for quasars at $z \sim 4$. We can therefore calculate the space densities of *all quasars* at this z , as a function of $M_{AB}(1450\text{\AA})$, simply by dividing the corresponding space densities of the radio-loud quasars by G_{RL} e.g.:

$$\rho(M_{AB}(1450\text{\AA}) \leq -26^m9) = (7.4 \pm 2.6) \text{ Gpc}^{-3}$$

and

$$\rho(M_{AB}(1450\text{\AA}) \leq -27^m5) = (3.8 \pm 1.7) \text{ Gpc}^{-3}$$

These numbers are consistent with the results of Fan et al. (2001b) at $z = 4.3$ (Fig. 4). Their maximum-likelihood luminosity function gives a density of $4.1^{+3.8}_{-2.1} \text{ Gpc}^{-3}$ at $M_{AB}(1450\text{\AA}) \leq -26.9^m$.

In Fig. 4 we compare our cumulative quasar luminosity function with that from 2dF at $z \approx 2$ (Boyle et al. 2000). We use the actual data (their Fig. 4), rather than their pure-luminosity-evolution model fit, which tends to overestimate the space densities at the bright end (both are shown in Fig. 4). In order to make this comparison we have converted their M_B to $M_{AB}(1450\text{\AA})$ assuming $M_B = M_{AB}(1450\text{\AA}) + 1.2\alpha_{opt} + 0.12$ (Kennefick et al. 1995) and adopting $\alpha_{opt} = -0.5$ (Boyle et al. 2000). In estimating the errors we allow also for possible systematic differences of 0.1 in the magnitude scales. The 2dF space density at $z \approx 2$ is:

$$\rho(M_{AB}(1450\text{\AA}) \leq -26^m9) = (13.5 \pm 4) \text{ Gpc}^{-3}$$

Thus the ratio between the space densities, for $M_{AB}(1450\text{\AA}) < -26^m9$, at $z \approx 2$ and at $z \approx 4$ is:

$$\rho_{z2}/\rho_{z4} = 1.8 \pm 0.8$$

If we extrapolate the $z = 2$ luminosity function to brighter magnitudes we get:

$$\rho(M_{AB}(1450\text{\AA}) \leq -27^m5) \approx 2.0 \text{ Gpc}^{-3}$$

and the ratio is less than 1 at $M_{AB}(1450\text{\AA}) \leq -27^m5$.

⁵The definition of radio-loudness is rather arbitrary. We actually define as 'radio loud' any quasar detected in FIRST, i.e. $S_{1.4\text{GHz}} \geq 1 \text{ mJy}$, which corresponds to $\log P_{1.4\text{GHz}} \geq 25.7$ at the median redshift of the $z = 4$ sample. Gregg et al. (1996) use the very similar criterion $\log P_{1.4\text{GHz}} > 25.5$.




Fig. 3.— The plots of Radio and Optical luminosity vs z , for the Q_{z4RL} sample (left plots) and for the Q_{z2RL} sample (right plots). Filled symbols are for the Einstein – de Sitter cosmology, empty symbols are for the *concordance model* (see Sect. 3.2).




Fig. 4.— The cumulative luminosity function of optically-luminous quasars at $z \sim 4.2$, derived in Sect. 4.2 (filled dots, 2 sample error bars shown), compared with SDSS results for $z = 4.3$ (shaded area, $\pm 1\sigma$ confidence level). The solid curve is the cumulative luminosity function obtained by integrating the $2.0 < z < 2.3$ luminosity function from the 2dF survey (Boyle et al. 2000), the dashed curve is their global model best fit.

Had we used the pure-luminosity-evolution model the densities would have been larger by a factor ≈ 1.7 and ≈ 2.0 and the density ratios would have been ≈ 3.0 and ≈ 1 .

The change in space-density ratio with optical luminosity is evident in Fig. 4.

5. Discussion

We have shown that the space density of optically-luminous quasars, $M_{AB}(1450\text{\AA}) < -26^m9$, declines moderately, if at all, from $z \approx 2$ to $z \approx 4$, contrasting sharply with the steep decline (factor ~ 10) found for samples including lower luminosity quasars (Shaver et al. 1996, 1999, Warren et al. 1994, Schmidt et al. 1995, Kennefick et al. 1995).

We stress that the quasars in our sample come from the bright tail of the optical luminosity function. The difference between our result and those previously reported is due to the different cosmological histories of quasars of different optical luminosity, the space density of the highly-luminous ones having peaked at earlier epochs than that of the less luminous ones.

This abundance of very bright quasars at $z \approx 4$ has already been noted by Fan et al. (2001b), who showed that their luminosity function at $z \simeq 4$ is flatter than that at lower redshifts (note that this is not readily-apparent in their Fig. 4, where the $z \approx 2$ 2dF-luminosity function is not reported correctly). Our $z = 4$ cumulative luminosity function is broadly consistent with that of Fan et al. (2001b) for SDSS, but extends it by about one decade in optical luminosity.

The relatively high space density of optically-luminous quasars at high redshift, and the consequent existence of super-massive black holes at very early stages of the evolution of the universe, yield non-trivial constraints on models for structure formation (e.g.: Haehnelt & Rees, 1993; Granato et al. 2001; Haiman & Loeb 2001).

6. Conclusions

We sought $z \geq 3.8$ quasars identified with FIRST radio sources $S_{1.4GHz} > 1$ mJy (irrespective of radio spectral index), i.e. 30 times fainter than previous searches, and built a sample of 13 optically-bright radio-loud quasars

($M_{AB}(1450\text{\AA}) \leq -26^m.9$, $\log P_{1.4\text{GHz}}(\text{W Hz}^{-1}) \geq 25.7$) with $3.8 < z < 4.5$. Our conclusions are:

- The space density of quasars meeting the above luminosity criteria is $(1.0 \pm 0.3) \text{ Gpc}^{-3}$.
- From a comparison sample at $z \approx 2$ we find that the space density of radio-loud quasars declines between $z \approx 2$ and $z \approx 4$ by a factor 1.9 ± 0.7 .
- We obtained an improved measurement of the radio-loud fraction of quasars in the luminosity range $\log P_{1.4}(\text{W Hz}^{-1}) > 25.7$: $G_{RL} = (13.4 \pm 3)\%$, nearly independent of optical luminosity, in the range $-25^m \gtrsim M_{AB}(1450\text{\AA}) \gtrsim -29^m$.
- Using the above value of G_{RL} , we calculate the space density of optically-selected quasars with $M_{AB}(1450\text{\AA}) < -26^m.9$, to be $(7.4 \pm 2.6) \text{ Gpc}^{-3}$. Using the 2dF quasar luminosity function, we find a ratio $\rho_{z2}/\rho_{z4} = 1.8 \pm 0.8$, consistent with that for the radio-loud quasars, but significantly different from the generally quoted ratio ~ 10 found for samples including lower-luminosity quasars.
- The moderate decline of quasar space density from $z = 2$ to $z = 4$ may halt altogether at the very bright end of the optical luminosity function, as, to some extent, is apparent in results from SDSS (Fan et al. 2001b).

7. Appendix : Absolute magnitude computation

We obtained the continuum absolute magnitudes of the quasars, $M_{AB}(1450\text{\AA})$ using:

$$M_{AB}(1450\text{\AA}) = E - 43.264 - 5\log(d_L(z)) - 2.5\log \int S_{\lambda}^{\text{Vega}}(\lambda) T(\lambda) d\lambda - 2.5\log(1+z) + 2.5\log \int \frac{S_{\lambda}^Q(\lambda/(1+z))}{S_{\lambda}^Q(1450\text{\AA})} T(\lambda) d\lambda \quad (1)$$

E is the APS POSS-I magnitude corrected for Galactic extinction. AB apparent magnitudes are defined by $m_{AB}(\lambda) = -2.5\log S_{\nu}(\lambda) - 48.60$, where S_{ν} is measured in $[\text{erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}]$ (Oke 1974). $S_{\lambda}^{\text{Vega}}$ in $[\text{erg cm}^{-2} \text{ s}^{-1} \text{\AA}^{-1}]$ was taken




Fig. 5.— The top panel shows the FBQS composite quasar spectrum (Brotherton et al. 2001), overplotted with a power law with spectral index $\alpha_{\nu} = -0.5$ (dotted curve). The bottom panel shows the quasar template (solid curve) in the observer's frame for $z = 4.2$, after correction for the Madau (1995) transmission. The POSS-I E response is shown as a dashed curve.

from Fukugita et al. (1996). S_{λ}^Q is based on the FBQS composite by Brotherton et al. (2001) and on the models of Madau (1995) for intervening absorption. First, an unabsorbed FBQS composite was created using the Madau model for $z = 2.5$. This was then corrected for intervening absorption at each redshift using linear interpolations of Madau's transmission curves for $z = 3.5$ and $z = 4.5$ (see Fig. 5 for an example). $T(\lambda)$ is the response function of the POSS-I E filter + plate (Evans 1989) and d_L is the luminosity distance in Mpc. The E magnitude of Vega, $E(\text{Vega}) = 0^m.05$, was obtained from the relation $E - R$ vs $V - R$ by Humphreys et al. (1991), using $V(\text{Vega})$ and




Fig. 6.— The K -correction between apparent magnitude $m_{AB}[1450(1+z)]$ and apparent E magnitude (POSS-I plate+filter response). The solid line corresponds to $\text{Ly}\alpha + \text{NV}$ equivalent width 70 Å, the dotted line to 15 Å.

$R(\text{Vega})$ from Fukugita et al. (1995).

The combined $\text{Ly}\alpha + \text{NV}$ rest-frame equivalent width (EW) of the FBQS composite is 70 Å at $z = 4.2$, close to the mean, 75 Å (rms ~ 30 Å), for our $z = 4$ sample. The $\text{Ly } \alpha$ line of our quasar 0918+0636 is heavily extinguished, $\text{EW}(\text{Ly } \alpha + \text{NV}) \leq 15$ Å. For this quasar the emission-line contour of the template was scaled to $\text{EW} = 15$ Å.

The relation between $m_{AB}[1450(1+z)]$ and E is shown as a function of redshift for $3.8 < z < 4.5$ in Fig. 6 for the two values of EW. E magnitude decreases with increasing strength of the emission line, particularly at $z \sim 4.2$, when $\text{Ly}\alpha + \text{NV}$ fall in the middle of the E band.

We thank Prof. G. Zamorani for useful discussion and criticism. RC and JIGS acknowledge financial support from DGES project PB98-0409-C02-02 and MCYT project AYA 2002-03326. KHM was supported by a Marie-Curie Fellowship of the European Commission. JH acknowledges funding via a PPARC studentship. This research has made use of the APS Catalog of POSS-I, which is supported by NASA and the University of Minnesota; and of the NASA/IPAC Extragalactic Database (NED) which is operated by the

Jet Propulsion Laboratory, California Institute of Technology under contract with NASA. We thank an anonymous referee for his comments and suggestions which helped in improving the presentation of the paper.

REFERENCES

- Becker R.H., White R.L., Helfand D.J. 1995, *ApJ* 450, 559
- Becker R.H., White R.L., Gregg M.D. et al., 2001a, *ApJS* 135, 227
- Becker R.H., Fan X., White R.L. et al., 2001b, *AJ* 122, 2850
- Benn C.R., Vigotti M., Pedani M. et al., 2001, *MNRAS* 329, 221
- Bertin E., Arnouts S., 1996 *A&A* 117, 393
- Boyle B.J., Shanks T., Croom S.M., Smith R.J., Miller L., Loaring N., Heymans C., 2000, *MNRAS* 317, 1014.
- Brotherton et al. 2001, *ApJ* 546, 775
- Djorgovski G. (October 2001 release), URL: <http://www.astro.caltech.edu/~george/z4.qsos>
- Dunlop J.S., Peacock J.A., 1990, *MNRAS* 247, 19
- Evans D.W., 1989, *AAS* 78, 249
- Fan X., Narayanan V.K., Lupton R.H. et al., 2001a, *AJ* 122, 2833
- Fan X., Strauss M.A., Schneider D.P. et al., 2001b, *AJ* 121, 54
- Fan X. et al., 2001c, *AJ* 121, 31
- Fukugita M., Shimasaku K., Ichikawa T., 1995, *PASP* 107, 945
- Fukujta et al. 1996, *AJ* 111, 1748
- Granato G.L., Silva L., Monaco P., Panuzzo P., Salucci P., De Zotti G., Danese L., 2001, *MNRAS* 324, 757
- Gregg et al., 1996, *AJ* 107, 1270
- Gregory P.C. et al., 1996, *ApJS* 103, 427
- Haiman Z., Loeb A., 2001, *ApJ* 552, 459

- Haehnelt M.G., Rees M.J., 1993, MNRAS 263, 168
- Hook I.M. et al., 1998a, MNRAS 297, 1115
- Hook I.M., Shaver P.A., McMahon R.G., 1998b, ASP Conf. Ser. 146: The Young Universe: Galaxy Formation and Evolution at Intermediate and High Redshift, S. D’Odorico, A. Fontana, E. Giallongo eds., p. 17
- Hook I.M. et al., 2002, A&A 391, 509
- Humphreys R.M. et al., 1991, AJ 102, 395
- Irwin M., McMahon R., Hazard C., 1991, in The Space Distribution of Quasars, edited by Crampton, ASPCS 21, 117.
- Ivezic et al., 2002, AJ 124, 2364
- Kennefick et al. 1995, AJ 110, 2563.
- Jarvis M.J., Rawlings S., 2000, MNRAS 319, 121
- Jarvis M.J., Rawlings S., Willott C.J., Blundell K.M., Eales S., Lacy M., 2001, MNRAS 327, 907
- Madau P. 1995, ApJS 441, 18
- Miyaji T., Hasinger G., Schmidt M., 2000, A&A 353, 25
- McMahon R. et al., 2002, ApJS 143, 1
- McMahon R., Irwin M.J., 1992, in Digitized Optical Sky Surveys, eds. H.T. McGillivray, E.B. Thomson (Dordrecht, Kluwer), p. 417
- Oke J.B. 1974, ApJS 27, 21
- Prandoni I. et al., 2001, A&A 365, 392
- Peacock J.A., 1985, MNRAS 217, 601
- Pennington R.L., Humphreys M., Odewahn C. et al., 1993, PASP 105, 521
- Schmidt M., Schneider D.P., Gunn J.E., 1995, AJ 110, 68
- Shaver P.A., Hook I.M., Jackson C.A., Wall J.V., Kellermann K.I., 1999, ASP Conf. Ser. 156: Highly Redshifted Radio Lines, C.L. Carrilli et al. eds., p. 163
- Shaver P.A., Wall J.V., Kellermann K.I., Jackson C.A., Hawkins M.R.S., 1996, Nature 384, 439
- Snellen I.A.G., McMahon R.G., Dennett-Thorpe J., Jackson N., Mack K.-H., Xanthopoulos E., 2001, MNRAS 325, 1167
- Stern D., Djorgovski S.G., Perley R.A., de Carvalho R.R., Wall J.V., 2000, AJ 119, 1526
- Vigotti M. et al., 1997, A&ASS 123, 1
- Warren S.J., Hewett P.C., Osmer P.S., 1994, ApJ 421, 412
- White R.L., Becker R.H., Helfand D.J., Gregg M.D. et al., 1997, ApJ 475, 479
- White R.L., Becker R.H., Gregg M.D. et al., 2000, ApJS 126, 133